

Math 4707: Integer partitions

Reminders: • HW#1 has been graded, visible on Canvas.

If you have questions/issues w/ grading, e-mail me.

- HW#2 is due this Wed., 2/17.
- Midterm #1 will be posted soon, is due Wed. of next week, 2/24.

We're almost done with **enumerative combinatorics**, our first major topic. Next class we will start with **graph theory**, the next major topic. Today we will discuss one last popular topic in enumerative combinatorics: **the study of (integer) partitions**.

Compositions

Before we discuss partitions, let's talk about **(integer) compositions**, which are closely related.

Defn A composition of n into k parts is a way of writing n as a sum of k positive integers:

$$n = a_1 + a_2 + \cdots + a_k.$$

e.g. $13 = 5+1+5+2$ is a composition of 13 into 4 parts.

$13 = 2+1+5+5$ is a different one.
order matters!

Q: how many comp. of n into k parts are there?

If we didn't have the requirement that $a_i \geq 1$, i.e., if we allowed $a_i = 0$ ('weak compositions')
then this is the same as the 'giving n pennies
to k children' problem, which had the formula

$$\binom{n+k-1}{k-1} \quad \text{via stars + bars.}$$

For requirement $a_i \geq 1$, easy fix: first give 1 penny to each kid, then distribute $n-k$ remaining pennies in $\binom{(n-k)+k-1}{k-1} = \binom{n-1}{k-1}$ ways.

\Rightarrow Thm # compositions of n into k parts
 $= \binom{n-1}{k-1}$

Cor # comp. of n into any number of parts
 $= 2^{n-1}$

Pf: Do sum $\sum_{k=1}^n \binom{n-1}{k-1} = \sum_{k=0}^{n-1} \binom{n-1}{k} = 2^{n-1}$ ✓

Note: Another direct way to see 2^{n-1} via stars+bars is:

$$n=5 \quad * * \uparrow * \uparrow * \uparrow \quad \text{e.g. } * | * * * * | * = 1+3+1$$

bar here? Y/N?

Partitions

A partition of n is like a composition, except that order doesn't matter:

$$1+5+5+2 \text{ same as } 2+1+5+5$$

Def'n A partition of n into k parts is a way of writing $n = \lambda_1 + \lambda_2 + \dots + \lambda_k$ w/ positive integers s.t. $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$.

↑
this convention chooses one fixed ordering
for any $\lambda_1, \dots, \lambda_k$.

e.g. We'd write above as $5+5+2+1$

Note Often write partition as **tuple** $\lambda = (\lambda_1, \dots, \lambda_k)$.

e.g.) $\lambda = (5, 5, 2, 1)$.

Let $p(n) := \# \text{ partitions of } n$.

n	partitions	$p(n)$
1	1	1
2	2, 1+1	2
3	3, 21, 111	3
4	4, 31, 22, 211, 1111	5
5	5, 41, 32, 311, 221, 2111, 11111	7
6	???	11


Q: What is $p(n)$?

A: Much, much harder to give formula for $p(n)$ than case of compositions.

Even if we won't easily be able to count them, let's think a little more about partitions ...

\exists a very nice graphical representation of a partition, called its Young diagram:

$$4+4+2+1 \Leftrightarrow$$




$$= \lambda$$

We see a new symmetry from Young diagram:

λ partition, its conjugate λ^t has transposed Young diagram:

$$4+3+2+2 \Leftrightarrow$$




$$= \lambda^t$$

Prop: # partitions of n into k parts

= # partitions w/ largest part = k .
of n


Pf: ???



Can we say anything about self-conjugate partitions (i.e., equal to own conjugate)?

Thm. # Self-conjugate partitions of n
= # partitions of n into distinct, odd parts.

Pf. Look at this picture:



any self-conjugate partition
can be decomposed into "elbows"
like this \square

We'll see another bijection between two classes
of partitions in a moment, but first...


Generating functions

In the past two classes we've seen that
generating functions are very powerful tools
to help understand sequences, and so what
about using them for partitions.

$$\text{Prop. } \sum_{n \geq 0} p(n) q^n = \prod_{j=1}^{\infty} \frac{1}{1-q^j}.$$

Pf. $\frac{1}{1-x} \cdot \frac{1}{1-x^2} \cdot \frac{1}{1-x^3} \cdots \cdot \frac{1}{1-x^k} \cdots =$

$$(1+x+x^2+\dots)(1+x^2+x^4+\dots)(1+x^3+x^6+\dots)\cdots(1+x^k+\dots)$$



Every choice of a term from each factor creates a unique partition, w/ exponent of $n = \text{size}$.

Note: We saw before that for rational generating functions $\frac{P(x)}{Q(x)} = \sum a_n x^n$, rate of growth of a_n has to do w/ 'singularities', i.e. roots of $Q(x)$.

Singularities of $\prod_{j=1}^{\infty} \frac{1}{1-q^j}$ are much harder to analyze

St: II, w/ (a lot of) work, idea leads to

Thm (Hardy-Ramanujan)

$$p(n) \sim \frac{1}{4n\sqrt{3}} e^{\pi\sqrt{2n/3}} \quad \text{as } n \rightarrow \infty.$$

What about a more down-to-earth use of g.f.'s for partitions? Can prove more identities ...

Let $O_n = \#$ partitions of n into odd parts
 $d_n = \#$ partitions of n into distinct parts

e.g.

n	odd	distinct	O_n	d_n
1	1	1	1	1
2	11	2	1	1
3	3, 111	3, 21	2	2
4	31, 1111	4, 31	2	2
5	5, 311, 11111	5, 41, 32	3	3

$$O(x) := \sum_{n \geq 0} O_n x^n = \frac{1}{1-x} \cdot \frac{1}{1-x^3} \cdot \frac{1}{1-x^5} \cdots$$

why?

$$D(x) := \sum_{n \geq 0} d_n x^n = (1+x)(1+x^2)(1+x^3) \cdots$$

∅ □ ∅ □ ∅ □ etc.

Then,

$$D(x) \cdot \frac{(1-x)}{(1-x)} \cdot \frac{(1-x^2)}{(1-x^2)} \cdot \frac{(1-x^3)}{(1-x^3)} \cdot \frac{(1-x^4)}{(1-x^4)} \cdots$$

$$= (1+x)(1+x^2)(1+x^3) \cdots \frac{(1-x)}{(1-x)} \frac{(1-x^2)}{(1-x^2)} \frac{(1-x^3)}{(1-x^3)} \cdots$$

$$= \frac{(1+x)(1-x)}{(1-x)} \cdot \frac{(1+x^2)(1-x^2)}{(1-x^2)} \cdot \frac{(1+x^3)(1-x^3)}{(1-x^3)} \cdots \frac{\cdots}{(1-x^n)}$$

recall $(1-a)(1+a) = (1-a^2)$

$$\cdots = \frac{1}{1-x} \cdot \frac{1}{1-x^3} \cdot \frac{1}{1-x^5} \cdots = O(x).$$

Cor. $d_n = o_n$ for all n . Pf: extract x^n coeff. \Rightarrow

MAGIC !!!

Note: \exists bijection $O(n) \xrightarrow{\sim} D(n)$ using
binary representation!

e.g. $\overset{n}{\overbrace{26}} = 5 + 5 + \underbrace{5 + 3 + 3 + 1 + 1 + 1 + 1 + 1}_{\text{odd parts}}$

$$= 3(5) + 2(3) + 5(1)$$

$\overset{\text{binary}}{\overbrace{26}} = (2^1 + 2^0)5 + (2^1)(3) + (2^2 + 2^0)(1)$

$$= 10 + 5 + 6 + 4 + 1$$

$\overset{\text{distinct parts!}}{\overbrace{10 + 5 + 6 + 4 + 1}}$

^{think abt. later...}

There are many more great theorems about partitions, like :

- Euler's pentagonal number theorem,
 - Ramanujan's congruences, . . .
- et cetera.

But now let's take a 5 min. break
and when we come back
we'll do the worksheet on
partitions in breakout groups ...