

Sign-reversing involutions + identities involving signs

Some identities w/ \pm signs can be proven like this:

Prop. Given a set X with a sign function $\text{sgn}: X \rightarrow \{\pm 1\}$
a weight function $\text{wt}: X \rightarrow \mathbb{R}$

and a sign-reversing, weight-preserving, involution
 $(\text{sgn}(\tau(x)) = -\text{sgn}(x)) \quad (\text{wt}(\tau(x)) = \text{wt}(x)) \quad (\tau^2 = \text{id})$

$$\text{then } \sum_{x \in X} \text{sgn}(x) \cdot \text{wt}(x) = \sum_{x \in X^2} \text{sgn}(x) \cdot \text{wt}(x)$$

$$= \sum_{x \in X : \tau(x) = x} \text{sgn}(x) \cdot \text{wt}(x)$$

Proof:

$$X = \begin{array}{|c|c|} \hline x^+ & x^- \\ \hline \text{cancel} & \\ \hline x & \tau(x) \\ \hline \end{array} \quad X \times X^2$$

$$\text{sgn}(x) \cdot \text{wt}(x) + \text{sgn}(\tau(x)) \cdot \text{wt}(\tau(x)) = 0$$

$$- \text{sgn}(x) \quad \text{wt}(x)$$

for all $x \in X \setminus X^2$

only this left \square

Examples

$$(1) \text{(Warm-up)} \quad \sum_{k=0}^n \binom{n}{k} (-1)^k = 0 \text{ for } n \geq 1$$

$$\sum_{\substack{\#S \\ \text{subsets } S \subseteq [n]}} (-1)^{\#S}$$

$$\text{sgn}: X = 2^{[n]} \rightarrow \{\pm 1\}$$

$$S \mapsto (-1)^{\#S}$$

$$\text{wt}: X = 2^{[n]} \rightarrow \mathbb{Z}$$

$$S \mapsto 1$$


$$\tau: X \rightarrow X$$

$$S \mapsto \begin{cases} S - \{1\} & \text{if } 1 \in S \\ S \cup \{1\} & \text{if } 1 \notin S \end{cases}$$

is sign-reversing,
weight-preserving
with no fixed points.

Rmk: This was key identity in pf. of P.I.E. \square

Pentagonal numbers:



② Recall Jhm (Euler's "Pentagonal Number Theorem")

$$\prod_{j \geq 1} (1 - q^j) = 1 + \sum_{n \geq 1} (-1)^n \left(q^{\frac{n(3n-1)}{2}} + q^{\frac{n(3n+1)}{2}} \right)$$

$$= 1 - q - q^2 + q^5 + q^7 - q^{12} - q^{15} + \dots$$

Franklin's (1881) proof of Euler's P.N.T.


$$\text{LHS} = (1-q)(1-q^2)\dots = \sum_{\lambda} \frac{\text{sgn}(\lambda)}{(-1)^{\ell(\lambda)}} \cdot \frac{q^{|\lambda|}}{|\lambda|!}$$

$\lambda :=$ partitions
 $\lambda = \lambda_1 > \lambda_2 > \dots > \lambda_r > 0$
 into distinct parts

$$\text{RHS} = 1 - q - q^2 + q^5 + q^7 - q^{12} - q^{15} + \dots$$

The diagram illustrates the expansion of the RHS series using geometric shapes. It shows three columns of rectangles. The first column, labeled n=1, contains one rectangle divided into four quadrants. The second column, labeled n=2, contains two rectangles, each divided into four quadrants. The third column, labeled n=3, contains three rectangles, each divided into four quadrants. The pattern indicates that the number of rectangles increases by one in each subsequent column, and each rectangle is divided into four quadrants.

Franklin defined $\gamma: X := \{\lambda \text{ w/ distinct parts}\} \rightarrow X$ by comparing
 • Smallest part and • longest initial run $\lambda_1, \lambda_1-1, \lambda_1-2, \dots$
 and moving the smaller one onto the bigger one:



When one can do this, check $\varphi^2 = \text{id}$, $\ell(\varphi(\lambda)) = \ell(\lambda) \pm 1$, $(\lambda) = |\varphi(\lambda)|$
 One cannot do this if:

- smallest part +
run have the same
size and overlap

$$n=3, \lambda_1 = \frac{3a(n)}{2}$$

- or runs
 - 1 smaller t
 - 1) they overlap

$$N = \frac{3n(n+1)}{2}$$

So χ implies only these shapes contribute to LHS \Rightarrow LHS = RHS.

11/5

(3) Theorem (Kirchoff's Matrix-Tree Theorem)


The number of spanning trees in a multigraph $G = \langle V, E \rangle$
for any i is $\det(\tilde{L}(G)^{i,i})$, where $\tilde{A}^{i,i}$ means A w/ row+column i removed,
(multiple/parallel edges allowed)

and $L(G)$ is the $n \times n$ Laplacian matrix of G :

$$L(G)_{v,w} := \begin{cases} \deg(v) & \text{if } v=w \\ -\# \text{edges from } v \text{ to } w & \text{if } v \neq w \end{cases}$$

Note! A spanning tree T of G is a subgraph of G that's a tree and which contains all the vertices V .

Example $G = \langle \{a, b, c, d\}, E \rangle$ has 5 spanning trees:



$$\text{And } L(G) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & -1 \\ 3 & -1 & 2 \end{bmatrix}$$

$$\text{So } \det(L(G)^{1,1}) = \det([3]) = 6 - 1 = 5 \checkmark$$

$$\text{and } \det(L(G)^{3,3}) = \det([-2 3]) = 9 - 4 = 5 \checkmark$$

Example Recall Cayley's formula n^{n-2} for # of (labeled) trees on n vertices. These are the ...

Spanning trees of the complete graph K_n on $[n]$.

e.g. $n=5$  $\overline{L(K_5)}^{5,5} = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & n-1 & \dots & \dots & n \\ 3 & \dots & n-1 & \dots & \dots \\ 4 & \dots & \dots & n-1 & \dots \\ 5 & \dots & \dots & \dots & n-1 \end{bmatrix} = n \underbrace{\begin{bmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{bmatrix}}_{(n-1) \times (n-1) \text{ identity matrix}} - \underbrace{\begin{bmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{bmatrix}}_{\text{all } 1's \text{ matrix}} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \end{bmatrix}$

What are eigenvalues of $\mathbb{1}_{n-1}$? It has rank = 1, so $(n-2)$ eigenvalues = 0
Also $\mathbb{1}_{n-1} \cdot \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} = (n-1) \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$, so one eigenvalue = $(n-1)$.


$\mathbb{1}_{n-1}$ has eigenvalues $(0, 0, \dots, 0, n-1) \Rightarrow \overline{L(K)}^{n,n}$ has eigenvalues $(n, n, \dots, n, 1)$

$$\Rightarrow \det(\overline{L(K)}^{n,n}) = n^{n-2} \Rightarrow \text{Cayley's formula.} \checkmark$$

In fact, let's prove a weighted, directed version of Kirchoff!

Thm If $L = \begin{bmatrix} 1 & -a_{12} & -a_{13} & \dots & -a_{1n} \\ 2 & -a_{21} & 1 & -a_{23} & \dots & -a_{2n} \\ 3 & \vdots & \ddots & \ddots & \ddots & \vdots \\ n & -a_{n1} & \dots & \dots & \dots & 1 \end{bmatrix}$ has $L_{ij} = \begin{cases} -a_{ij} & \text{if } i \neq j \\ a_{i1} + a_{i2} + \dots + \hat{a_{ij}} + \dots + a_{in} & \text{if } i = j \end{cases}$

then $\det(\overline{L}^{K,K}) = \sum_{\substack{\text{arborescences} \\ \text{A directed toward } K}} \prod_{i \rightarrow j \text{ in } A} a_{ij}$, where a_{ij} are formal parameters.




Note: \Rightarrow Kirchoff by setting $a_{ij} = \# \text{edges } i \rightarrow j$ in G

$$L = \begin{bmatrix} 1 & 2 & 3 \\ 2 & a_{12}+a_{13} & -a_{12} & -a_{13} \\ 3 & -a_{21} & a_{21}+a_{23} & -a_{23} \\ 1 & -a_{31} & -a_{32} & a_{31}+a_{32} \end{bmatrix} \Rightarrow \det(\overline{L}^{3,3}) = \det \begin{bmatrix} a_{12}+a_{13} & -a_{12} & 0 \\ -a_{21} & a_{21}+a_{23} & 0 \\ 0 & 0 & a_{31}+a_{32} \end{bmatrix}$$

$$= (a_{12}+a_{13})(a_{21}+a_{23}) - (-a_{12})(-a_{21})$$

$$= a_{12}a_{23} + a_{12}a_{21} + a_{13}a_{21} + a_{13}a_{23} - a_{12}a_{21}$$

$$= a_{12}a_{23} + a_{13}a_{21} + a_{13}a_{23} \checkmark$$



11/8

Proof of Thm:

$$L = \begin{bmatrix} R_1 - a_{11} & -a_{12} & \cdots & -a_{1n} \\ -a_{21} & R_2 - a_{22} & \cdots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ -a_{n1} & \cdots & R_n - a_{nn} \end{bmatrix}$$

where $R_i = a_{i1} + a_{i2} + \cdots + a_{ii} + \cdots + a_{in}$

$$= \sum_{j=1}^n a_{ij}$$

$$= (R_i \cdot \text{sgn}(w) - a_{ij})_{\substack{i=1, \dots, n \\ j=1, \dots, n}}$$

$\Rightarrow \det(L^{n,n}) = \sum_{w \in S_{n-1}} \text{sgn}(w) \prod_{i=1}^{n-1} L_{i,w(i)}$

Recall: $\text{sgn}(w) = (-1)^{\text{inv}(w)}$

"Leibniz formula"

$$= \sum_{S \subseteq [n-1]} \prod_{i \in S} (R_i - a_{ii}) \sum_{w \in S_{[n-1] \setminus S}} \text{sgn}(w) \prod_{i \in [n-1] \setminus S} (-a_{ii}, w(i))$$

(fixed by w)

$$= \sum_{S \subseteq [n-1]} \sum_{T \subseteq S} \prod_{i \in T} R_i \prod_{i \in S \setminus T} (-a_{ii}) \sum_{w \in S_{[n-1] \setminus S}} \text{sgn}(w) \cdot \prod_{i \in [n-1] \setminus S} (-a_{ii}, w(i))$$

derangement

Important property of sign of permutations:
 $\text{sgn}(v \cdot w) = \text{sgn}(v) \cdot \text{sgn}(w)$

$$= \sum_{T \subseteq [n-1]} \prod_{i \in T} (a_{i1} + a_{i2} + \cdots + a_{in}) \cdot \sum_{w \in S_{[n-1] \setminus T}} \text{sgn}(w) \prod_{i \in [n-1] \setminus T} (-a_{ii}, w(i))$$

$$= \sum_{f: T \rightarrow [n]} \prod_{i \in T} a_{i, f(i)}$$

$$= \sum_{\substack{(T, f, w) \\ T \subseteq [n-1] \\ f: T \rightarrow [n] \\ w \in S_{[n-1] \setminus T}}} (-1)^{\#([n-1] \setminus T)} \cdot \frac{\text{sgn}(w)}{\text{sgn}(x)} \cdot \prod_{i \in T} a_{i, f(i)} \prod_{i \in [n-1] \setminus T} a_{i, w(i)}$$

$$\chi := \sum_{\substack{(T, f, w) \\ T \subseteq [n-1] \\ f: T \rightarrow [n] \\ w \in S_{[n-1] \setminus T}}} (-1)^{\#([n-1] \setminus T)} \cdot \frac{\text{sgn}(w)}{\text{sgn}(x)} \cdot \prod_{i \in T} a_{i, f(i)} \prod_{i \in [n-1] \setminus T} a_{i, w(i)}$$

We will evaluate this signed, weighted sum using a sign-reversing involution ...

Picture of (T, f, ω) :



We can define an involution
 $\chi: X \rightarrow X$
 that swaps the cycle containing
 the smallest index $i \in [n-1]$
 from w to f or back from f to w !


Check that χ

- is an involution (clear)
- is wt-preserving (preserves arcs)
- is sign-reversing (sgn of a k -cycle is $(-1)^{k+1}$)

What are the fixed points X^χ ?

No cycles \Rightarrow $[n-1] \setminus T$ is empty, i.e., $T = [n-1]$
 in w or f and $f: [n-1] \rightarrow [n]$ has no cycles

$$T = [n-1]$$



(easy) Lemma.

This forces f to be an arborescence directed toward n (and conversely, any arborescence is such an f).

Hence, $\det(L^{n,n}) = \sum_{\substack{\text{arborescences} \\ f \text{ of } [n] \text{ directed} \\ \text{toward } n}} \prod_{i \in [n-1]} a_{i, f(i)}.$

11/10

The transfer matrix method (Stanley § 4.7, Ardila § 3.1.2)

Another tool from linear algebra for counting walks in graphs

Thm Let G be a graph w/ vertex set $V = [n]$, and let

$A_G = (a_{ij})_{i,j=1,\dots,n}^{i=1,\dots,n}$ be its adjacency matrix: $a_{ij} = \# \text{edges from } i \text{ to } j$.

Then (a) # of walks of length ℓ = $(A_G^\ell)_{i,j}$ for all $\ell \geq 0$.

(b) # of closed walks of length ℓ
 $i = i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_\ell = i$ (for all i) = $\lambda_1^\ell + \lambda_2^\ell + \dots + \lambda_n^\ell$,
where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of A_G .

Pf: (a) is just definition of matrix multiplication:

$$(A_G^\ell)_{i,j} = \sum_{i_1=1}^n \sum_{i_2=1}^n \dots \sum_{i_{\ell-1}=1}^n a_{i,i_1} a_{i_1,i_2} \dots a_{i_{\ell-1},j} = \text{LHS of (a)}.$$

For (b), from (a) it follows that

$$\# \text{closed walks of length } \ell = \sum_{i=1}^n (A_G^\ell)_{i,i} = \text{trace}(A_G^\ell).$$

Since A_G is real + symmetric, it can be diagonalized,

i.e., $\exists P$ s.t. $P A_G P^{-1} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 & \dots & 0 \\ & & \ddots & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$. Thus,

$$\begin{aligned} \text{trace}(A_G^\ell) &= \text{trace}((P A_G P^{-1})^\ell) = \text{trace}((P^T \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 & \dots & 0 \\ & & \ddots & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix} P)^\ell) \\ &= \text{trace}(P^T \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 & \dots & 0 \\ & & \ddots & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}^\ell P) = \text{trace}(\begin{pmatrix} \lambda_1^\ell & 0 \\ 0 & \lambda_2^\ell & \dots & 0 \\ & & \ddots & \\ 0 & 0 & \dots & \lambda_n^\ell \end{pmatrix}) = \lambda_1^\ell + \dots + \lambda_n^\ell. \end{aligned}$$

Recall:
 $\text{tr}(AB) = \text{tr}(BA)$

Example Let $f(n, k) = \# \text{proper vertex-colorings of } C_n$ cycle graph
w/ k -colors.
(no adjacent vertices w/
Same color)

$$\text{e.g. } n=2 \quad f(2, k) = k(k-1)$$

$\frac{(k-1)(k-2)}{2} = \frac{k(k-1)}{2}$


color 1 first in k ways color 2 second in $k-1$ ways

$$\begin{aligned} n=3 \quad & f(3, k) = k(k-1)(k-2) \\ & \text{color 1} \quad \text{color 2} \quad \text{color 3} \end{aligned}$$

$$n=4 \quad f(4, k) = \underbrace{k(k-1)(k-2)(k-3)}_{\substack{2+4 \text{ have different} \\ \text{colors}}} + \underbrace{k(k-1)(k-1)}_{\substack{2+4 \text{ have} \\ \text{same color}}}^3$$

color 1 color 2 color 4 color 3 color 1 color 2+4
 ↓ ↓ ↓ ↓ ↓ ↓ ↓
 4 1 2 3 4 1 2+4
 3 2 4 1 3 2 Same

Note: $\{$ proper k -colorings of $C_n\} \leftrightarrow \{$ closed walks of length n in complete graph $K_k\}$



So taking $A_{K_k} = \begin{bmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \cdots & 0 \end{bmatrix} = I_k - J_k$

which has eigenvalues $(\lambda_1, \dots, \lambda_k) = (k-1, -1, -1, \dots, -1)$
 (since we saw earlier that I_k has eigen's $(k, 0, 0, \dots, 0)$)

we find that $f(n, k) = \lambda_1^n + \dots + \lambda_k^n$

$$\begin{aligned} &= (k-1)^n + (-1)^n + \dots + (-1)^n \\ &= (k-1)^n + (k-1)(-1)^n \\ &= (k-1)((k-1)^{n-1} + (-1)^n). \end{aligned}$$

e.g. $f(2, k) = (k-1)(k-1+1) = (k-1)k$
 $f(3, k) = (k-1)((k-1)^2 + 1) = (k-1)(k^2 - 2k + 1)$
 $f(4, k) = (k-1)((k-1)^3 + 1) = (k-1)(k^3 - 3k^2 + 3k + 1)$